arXiv:1212.0844v2 [cond-mat.stat-mech] 12 Jan 2013

Duration of violation of the second law of thermodynamics along a single trajectory in phase space

Reinaldo García-García* and Daniel Domínguez
Centro Atómico Bariloche and Instituto Balseiro, 8400 S. C. de Bariloche, Argentina

We study the fraction of time that a system spends violating the second law of thermodynamics within a single trajectory in the phase space under driving at a constant rate. We denominate this quantity as *violation ratio*, and we denote it by $\mathcal{R}_-(N,t) = \tau_-(N,t)/t$, where $\tau_-(N,t)$ is the amount of time the system has violated the second law up to time t in a system with N particles. We develop general arguments to predict that for generic ergodic systems its mean value scales, in the large-t - large-N limit, as $\langle \mathcal{R}_-(N,t)\rangle \sim (t\sqrt{N})^{-1}$. We determine, for a simple system driven at a constant rate, the precise asymptotic behaviour of the mean violation ratio, both, analytically and numerically, showing a good agreement with our prediction. Finally, we briefly discuss how this simple scaling may break down in the vicinity of a phase transition.

PACS numbers: 05.40.-a,05.70.Ln

Fluctuations in small systems induce violations of the second law of thermodynamics for arbitrary external protocols at individual trajectories. This fact is well characterized, in an ensemble-based picture, by the celebrated fluctuation theorems [1–6]. In this picture, the relative probability weight of those trajectories not violating the second law with respect to those violating it, is computed. This relative weight exhibits a simple form

$$\frac{P(X)}{P^{\mathrm{T}}(-X)} = \exp(X), \tag{1}$$

where T represents a transformation, usually time-reversal, or a transformation to a *dual* dynamics (see [7–9] for a simple definition of the dual dynamics), and X represents a trajectory-dependent thermodinamic quantity, as work, heat, or more generally, different forms of trajectory-dependent entropy production, possessing the symmetry $X^{\mathrm{T}} = -X$.

In the last few years, these relations have been extended to non-Markovian dynamics as well [10–16], and a lot of experimental work has been done, on one side, in order to test their validity in given systems, and on the other side, in order to use them, for example, to reconstruct the free energy landscape of small systems (see for instance [18–22]).

Eq.(1), which is commonly denominated a detailed fluctuation theorem (DFT), leads to the relation $\langle \exp(-X)\rangle = 1$, also known as an integral fluctuation theorem (IFT). From the IFT, the second law of thermodynamics can be immediately derived as $\langle X\rangle \geq 0$ whatever the protocol and the duration of the process are. In very large systems the fluctuations are naturally supressed and $\langle X\rangle \approx X$. Then, for any realization of any process one has $X \geq 0$. However, in small systems, fluctuations play an important role and realizations are possible such that $X < 0$ for not too large times. As previously stated, fluctuation theorems (FTs) express that these "second-law-violating" trajectories have an exponentially small weigth related to trajectories being compatible with the second law of thermodynamics.

One may wonder, however, if it is possible to characterize the way a system violates the second law from a different perspective. For instance, one may formulate the questions: Within a particular trajectory in the phase space, what amount of time (or what fraction of time) the system spends in the "violation sector" defined by $X < 0$? How does this fraction vanishes for large times and in the thermodynamic limit? In order to address these issues, we need to study the zero-crossings properties of the stochastic process $X(t)$. In particular, we should note that the amount of time, up to time t, that a process spends in the semi-space $X(t) < 0$ (or $X(t) > 0$), is called by the comunity of stochastic processes physicists as a "residence time". The probability density function (PDF) for this quantity is closely related to persistence, e.g. the probability that the process does not cross the line $X = 0$ up to time t [23]. This quantity have been studied in many different contexts. For example, in [23, 24] it have been used to characterize the ordering dynamics of the Ising model (where the process $X(t)$ is just the global or local magnetization).

Generally speaking, the computation of the full PDF of the residence time for a given stochastic process is a complex task, even is the process is Markovian. In our case, we choose X to be an stochastic entropy production (see definitions below) which is non-Markovian even if the system performs Markov dynamics. Thus, an exact determination of the full PDF is out of order. We rather derive some results for the first moment of this PDF, the average violation ratio, defined as $\langle \mathcal{R}_-(N,t)\rangle = \langle \tau_-(N,t)\rangle/t$, where $\tau_-(N,t)$ is the amount of time, up to time t, that within a given trajectory for the system with N particles, $X(t)$ was negative.

This kind of study may be relevant in some practical situations. For typical processes, violations of the second law of thermodynamics occur at very short times (thus, introducing a typical time scale related, of course, to the mean time the system spends in the sector of negative

X). Then, typical operation times of devices intended to work avoiding these effects should be greater than the introduced time scale, thus, there is a well defined limit for miniaturization, which can be properly determined by means of this kind of study.

In this letter, we consider generic ergodic and extensive systems driven by a set of external, well controled parameters (which we denote by $\ell(t)$), varying at constant rate. For constant protocols, the system is able to reach a unique steady-state characterized by the PDF $p_{\mathrm{s}}(\mathcal{C};\ell) = \exp[-\phi(\mathcal{C};\ell)]$, where $\mathcal{C}$ denotes a configuration in the phase space. $\mathcal{C}$ may be a single variable, a vector, or a field as well. We discuss equilibrium and non-equilibrium steady-states (NESS) on the same footing. Furthermore, we assume that the system is initially prepared in the steady-state corresponding to $\ell(0) = \ell_0$. We introduce the process $X(t)$ as follows

$$X(t) = \int_0^t ds \dot{\ell}(s) \partial_\ell \phi(\mathcal{C}(s); \ell(s)), \tag{2}$$

which depends on the particular trajectory of the system in the phase-space during its evolution. For equilibrium steady-states, this is no more than the total entropy production, proportional to the dissipated work [6] $\beta\mathcal{W}_{\mathrm{d}} = \beta\mathcal{W} - \beta\Delta F$, where $\mathcal{W}$ is the thermodynamical work introduced by Jarzynski [5], whereas F is the free energy. On the other hand, for NESS, this quantity corresponds to the functional of Hatano and Sasa [25], which is the correct choice in order to extend the second law of thermodynamics to NESS.

We start by advancing some fundamental results. For the generic kind of systems we are considering, the mean violation ratio exhibits, at large times, the simple behaviour $\langle \mathcal{R}_-(N,t)\rangle \sim \mathcal{A}(N)/t$, where $\mathcal{A}(N)$ is a model-dependent function of N which vanishes in the thermodynamic limit as $\mathcal{A}(N) \sim \mathcal{A}_\infty/\sqrt{N}$, with $\mathcal{A}_\infty$ a non-universal, model-dependent constant. We remark that, even if these results are very intuitive, we were not able to find them in the existing literature. We develop a general argument for ergodic systems in order to justify such a scaling law, and we test it in a simple Gaussian system both, analytically and numerically.

Let us begin by considering an ergodic system initially prepared in its steady state corresponding to a value of the parameters $\ell(0) = \ell_0$. We should note first that ergodicity has two important consecuences: *i*) the system satisfies the IFT (Jarzynski [5] or Hatano-Sasa [25])

$$\langle \exp(-X)\rangle = \int dX P(X,t) \exp(-X) = 1, \tag{3}$$

and *ii*) in the large-N limit, the probability density function of the extensive quantity X becomes a Gaussian

$$P(X,t) = \frac{1}{\sqrt{2\pi\Sigma_X(t)}} \exp\left[-\frac{\big(X - m_X(t)\big)^2}{2\Sigma_X(t)}\right], \tag{4}$$

where m_X is the mean value of the entropy production, while Σ_X coresponds to its variance. Non Gaussian corrections should be subleading in N as $N \to \infty$. However, some care has to be taken in order to take this limit when combined with the FTs as pointed out in [26], since the IFT $\langle \exp[-X]\rangle = 1$ is strongly influenced by rare trajectories contributing to the tails of the X-PDF, where the central limit theorem is weaker. We consider here the class of systems for which both, the fluctuation theorem and the Gaussian character of the fluctuations of X hold. Then in the large-N limit we have $\Sigma_X(t) = 2m_X(t)$, as can be easily seen by direct substitution of (4) in (3).

The second important point to note is that, for very short times, the mean entropy production per particle behaves as $\mu_X(t) = \frac{1}{2}\ddot{\mu}_X(0)t^2 + \mathcal{O}(t^3)$. This result follows from the fact that $m_X(0) = 0$ and $\dot{m}_X(0) = 0$. The first relation may be easily seen by using the definition of the entropy production $X(t)$. The second relation can be obtained by differenciating in (2), taking the average and having in mind that the system, at $t = 0$, is prepared in its steady-state.

Third, for generic processes the mean entropy production should increase with time. In particular, we consider protocols such that $m_X(t)$ is an strictly monotonous growing function (as it is the case for constant driving). Note that in this case, the system is able to violate the second law of thermodynamics only within a time sector where the mean entropy production is smaller than the standard deviation associated to the X-distribution. For larger times, as m_X increases, fluctuations are not strong enough to make X negative. We denote this characteristic time by $\tau_c(N)$. Thus, for $t > \tau_c(N)$, the average time the system spends violating the second law does not increase:

$$\langle \tau_-(t > \tau_c(N))\rangle = \mathrm{const} \le \tau_c(N). \tag{5}$$

This means that we can write $\langle \tau_-(t > \tau_c(N))\rangle = \mathcal{A}(N)$, with $\mathcal{A}(N) = \alpha(N)\tau_c(N)$, and $\alpha(N) \le 1$ a constant that depends in principle on the number of particles, the specific protocol and on temperature. We then have

$$\langle \mathcal{R}_-(N, t > \tau_c)\rangle = \mathcal{A}(N)/t \tag{6}$$

The previous argument explains the behaviour $\langle \mathcal{R}_-(N,t)\rangle \sim t^{-1}$, and also gives a precise meaning to the phrase "large times". Now, note that, as discussed before, the violation sector is defined by the relation $m_X(t) \lesssim \sqrt{2\Sigma_X(t)}$, which leads to $m_X(t) \lesssim 4$ using the derived relation between the mean and the variance of the X-PDF. Then $\tau_c(N)$ is determined by the relation $m_X(\tau_c) = N\mu_X(\tau_c) \approx 4$, leading to $\mu_X(\tau_c) \approx \frac{4}{N}$. Note that if N is large enough, then τ_c should be small. Thus, one can use the expansion of μ_X around $t = 0$, to obtain $\tau_c(N) \sim \frac{2\sqrt{2}}{\sqrt{N\ddot{\mu}_X(0)}}$, which means that we can write, to the leading order in t and N

$$\langle \mathcal{R}_-(N, t > \tau_c)\rangle \sim \frac{\mathcal{A}_\infty}{t\sqrt{N}}, \tag{7}$$

with $\mathcal{A}_\infty = \frac{2\sqrt{2}\alpha_\infty}{\sqrt{\ddot{\mu}_X(0)}}$, and $\alpha_\infty = \alpha(N \to \infty)$, which we assume finite. Equation (7) constitutes the main result of our work. In order to test it, we consider a discrete model of an elastic line where particles interact via a short-ranged elastic potential in the presence of periodic boundary conditions (PBC). Besides, the system is coupled to a parabolic trap. The center of the trap will be considered here as our external protocol. The Hamiltonian of the system is given, as usual, by

$$\mathcal{H}(\{u\}_i; \ell) = \sum_{i=1}^{N} (u_{i+1} - u_i)^2 + \frac{k}{2}\sum_{i=1}^{N} (u_i - \ell)^2, \tag{8}$$

where k denotes the strenght of the trap. It is simple to see that for the present system the free energy does not depend on ℓ, thus one has $\mathcal{W}_d = \mathcal{W}$. For the work, the relevant degree of freedom is the position of the center of mass of the chain

$$\mathcal{W}[u_{\rm cm}(s); \ell(s)] = kN\int_0^t ds\dot{\ell}(s)[\ell(s) - u_{\rm cm}(s)], \tag{9}$$

a result which follows by particularizing the general definition given in [5] to our case. The center of mass evolves as

$$\dot{u}_{\rm cm}(t) = k[\ell(t) - u_{\rm cm}(t)] + \eta(t), \tag{10}$$

where the effective noise $\eta(t)$ has zero mean and variance $\langle \eta(t)\eta(t')\rangle = \frac{2}{\beta N}\delta(t - t')$, if one assumes simple relaxational Langevin dynamics for the system at inverse bath temperature β. As can be seen, the center of mass moves like a particle interacting with a thermal bath equilibrated at the inverse temperature $\beta_{\rm eff}(N) = \beta N$. Then, for constant $\ell(t) = \ell$, the equilibrium distribution corresponds to

$$p_{\rm eq}(u_{\rm cm}, \ell) = \exp\left[-\beta\left(\frac{1}{2}Nk(u_{\rm cm} - \ell)^2 - \tilde{F}(\beta)\right)\right], \tag{11}$$

where $\tilde{F}(\beta)$ can be determined by normalizing $p_{\rm eq}(u_{\rm cm}, \ell)$. As our system is initially prepared in equilibrium, and by virtue of the previous discussion , we conclude that the work is a Gaussian process. We are interested in the case where the center of the parabola starts to move at a constant speed $\ell(t) = \ell_0 + vt$. Then, simple calcultions lead to the results $m_{\mathcal{W}}(t) = \langle \mathcal{W}(t)\rangle = N\mu_{\mathcal{W}}(t)$, and $\Sigma_{\mathcal{W}}(t) = \langle (\mathcal{W}(t) - m_{\mathcal{W}}(t))^2\rangle = \frac{2}{\beta}N\mu_{\mathcal{W}}(t)$, where $\mu_{\mathcal{W}}(t)$ is the average work per particle

$$\mu_{\mathcal{W}}(t) = v^2 t - \frac{v^2}{k}[1 - \exp(-kt)]. \tag{12}$$

With all this, the work PDF can be immediately written as a Gaussian of the form (4).

Now, we are in position to explicitly calculate the mean violation ratio

$$\langle \mathcal{R}_-(N, t)\rangle = \frac{1}{t}\int_0^t ds\langle \Theta(-\mathcal{W}(s))\rangle = \\ = \frac{1}{t}\int_0^t ds\int_{-\infty}^{0} dW P(W, s). \tag{13}$$

It is easy to see, from Eqs.(4) and (13), that for large t one has, to the leading order, that $\langle \mathcal{R}_-(N, t)\rangle \sim \mathcal{A}(N, v, \beta)/t$ for $t \to \infty$, with

$$\mathcal{A}(N, v, \beta) = \frac{1}{2}\int_0^\infty \mathrm{Erfc}\left(\frac{1}{2}\sqrt{\beta N\mu_{\mathcal{W}}(s)}\right)ds. \tag{14}$$

We are interested in the large N limit. In this case we can see, using (12), that to the leading order in $1/\sqrt{N}$ we have

$$\mathcal{A}(N, v, \beta) \approx \frac{1}{2\sqrt{N}}\int_0^\infty \mathrm{Erfc}\left(s\sqrt{\frac{\beta k v^2}{8}}\right)ds = \sqrt{\frac{2}{\pi\beta k v^2 N}}, \tag{15}$$

implying that

$$\langle \mathcal{R}_-(N, t)\rangle \sim \frac{\mathcal{A}_\infty}{t\sqrt{N}}; \quad t \to \infty, N \to \infty, \tag{16}$$

where $\mathcal{A}_\infty$ does not depend on t nor N.

We also perfomed some numerical simulations. Let us introduce the variable $y(t) = \ell(t) - u_{\rm cm}(t)$. In a time discretized picture, this new variable evolves as

$$y_n = y_{n-1} - (ky_{n-1} - v)\Delta t + \sqrt{\frac{2\Delta t}{\beta N}}\zeta_{n-1}, \tag{17}$$

where ζ_n is a random variable (for each n) such that $\langle \zeta_n\rangle = 0$, and $\langle \zeta_n^2\rangle = 1$. We use a time step $\Delta t = 0.005$. We first thermalize the system at $v_0 = 0$ during 50000 time steps and then we start to pull the parabola at a finite velocity v during a lapse of 20000 steps and compute the work at every time, and also the value of $\mathcal{R}_-(N, t)$. Each 200 steps (after the thermalization procedure) we save this quantity in order to compute its average. The work for each time step can be easily calculated as $\mathcal{W}_n = \mathcal{W}_{n-1} + kNv\Delta t y_{n-1}$.

Introducing the quantity f_n being one if the work is negative, and zero otherwise we have, using the general definition of $\mathcal{R}_-(N, t)$, that one can write $\mathcal{R}_-^{(n)} = \frac{n-1}{n}\mathcal{R}_-^{(n-1)} + \frac{1}{n}f_{n-1}$. If we introduce the scaling variable $\lambda = t\sqrt{N}$, we should have, for large t and N, that $\sqrt{\frac{\pi\beta k v^2}{2}}\langle \mathcal{R}_-(N, t)\rangle = G(\lambda)$. We set $\beta = 0.05$, $v = 1.0$, $k = 1.5$ and study different system sizes, say $N = \{2^4, 2^6, 2^8, 2^{10}, 2^{12}, 2^{14}, 2^{16}\}$. For large N, we find a very good agreement (see Fig. 1) with the expected scaling form $G(\lambda) = \lambda^{-1}$.



FIG. 1. (Color online) Scaling form of $\langle \mathcal{R}_-(N,t)\rangle$ in the elastic system. For large N, the law given by Eq. (16), with $\mathcal{A}_\infty = \sqrt{\frac{2}{\pi\beta k v^2}}$, is verified.

To finish, we discuss a simple case where the obtained scaling law may fail. First, let us write a more detailed expression for $\ddot{m}_X(0)$. Note that taking $\ell(t) = \ell_0 + vt$ we can write at short times

$$\langle \partial_\ell \phi(\mathcal{C}(t), \ell(t))\rangle \approx \langle \partial_\ell \phi(\mathcal{C}(t), \ell_0)\rangle + vt\langle \partial^2_\ell \phi(\mathcal{C}(t), \ell_0)\rangle_{\rm ss}, \tag{18}$$

where the double-s subscript means that the average is taken in the steady-state corresponding to $\ell = \ell_0$. On the other hand, to the same order, the normalization condition for the steady-state PDF leads to the relation $\langle \partial^2_\ell \phi(\mathcal{C}(t), \ell_0)\rangle_{\rm ss} = C_r(0)$, where the correlation function C_r is given by

$$C_r(t-s) = \langle \partial_\ell \phi(\mathcal{C}(t), \ell_0)\partial_\ell \phi(\mathcal{C}(s), \ell_0)\rangle_{\rm ss} \tag{19}$$

Also note that using the generalized fluctuation-dissipation relation (see, for instance, [27]) one can write

$$\langle \partial_\ell \phi(\mathcal{C}(t), \ell_0)\rangle = v\int_0^t ds s\chi(t-s), \tag{20}$$

with $\chi(t-s) = \partial_t C_r(t-s)$. With all this, using (2), we have

$$\dot{m}_X(t) \approx v^2 \int_0^t ds s\chi(t-s) + v^2 t C_r(0), \tag{21}$$

at short times, leading to the result $\ddot{m}_X(0) = v^2 C_r(0)$. Assume that our system has linear size L, being large, but finite, and that we are in the vicinity of a phase transition. Also assume that the correlation and response functions are associated to an order parameter. Then, the scaling form of the response function depends, for an infnite system, on the correlation length. However, if the system is finite the correlation length saturates, very close to the transition temperature, to L. In this case the response function in the frequency domain acquires the following scaling form [28]

$$\tilde{\chi}(\omega) = L^{2-\eta}F(\omega L^z) = N^{\frac{2-\eta}{d}}F_1(\omega N^{z/d}), \tag{22}$$

where $F(x)$ and $F_1(x)$ are scaling functions, z is the dynamic exponent, d is the spatial dimension and η corresponds to anomalous dimension exponent. From the previous analysis one sees that

$$\ddot{m}_X(0) = v^2 C_r(0) = v^2 \int_{-\infty}^{\infty} \frac{d\omega}{2i\pi\omega}\tilde{\chi}(\omega), \tag{23}$$

which leads to $\ddot{m}_X(0) = \mathcal{B}N^{\frac{2-\eta}{d}}$, where the constant $\mathcal{B} = v^2 \int_{-\infty}^{\infty} \frac{dx}{2i\pi x}F_1(x)$. Then we should have, if the general arguments we have developed before apply, that $\langle \mathcal{R}_-(N,t)\rangle \sim (tN^{\frac{2-\eta}{2d}})^{-1}$. It is worth noting that we must demand $2-\eta > 0$. This result is already different from the general law given by Eq. (7). Furthermore, it could even suggest that in this case the analysis leading to (7) is incorrect because fluctuations are so important that one can no longer assume that there is an small time τ_c such that for $t > \tau_c$ the system does not violate the second law, which implies that using the small t expansion for m_X is no longer correct. This fact could also affect the scaling form in the time variable. In other words, the previous theory breakes down since the typical relaxation time of the system diverges in the thermodynamic limit. Finally, note that in the mean-field case (where $\eta = 0$ for a scalar field) fluctuations are naturally unimportant. In this case the analysis leading to (7) is correct and one should have different scaling forms with N depending only on the spatial dimension in the vicinity of a phase transition ($\mathcal{R} \sim (tN^{1/d})^{-1}$), while far away from the phase transition (7) should hold.

In conclusion, we have introduced a general set of arguments to predict that for generic ergodic systems satisfying the fluctuation relations, the mean fraction of time the systems spends violating the second law of thermodynamics within a single trajectory scales, in the large-t - large-N limit, as $(t\sqrt{N})^{-1}$. This result could be useful for miniaturization purposes. We tested our theory in a simple system, both, analytically and numerically, finding a good agreement with the predicted scaling. We point out that non-ergodic systems, as well as systems where fluctuations are strong (as in the vicinity of a phase transition) may violate this simple scaling law. It would be interesting to compute the full distribution of the violation ratio to elucidate if some symmetry relation (reminiscent of the FTs) arises. On the other hand, a precise study of this distribution in the thermodynamic limit could give direct information about how the divergence of a relaxation time affects the second law of thermodynamics.

This work was supported by CNEA, CONICET (PIP11220090100051), and ANPCYT (PICT2011-1537).

---

* reinaldo.garcia@cab.cnea.gov.ar

[1] Denis J. Evans, E. G. D. Cohen and G. P. Morriss, Phys. Rev. Lett. **71**, 2401 (1993).

[2] G. Gallavotti and E. G. D. Cohen, Phys. Rev. Lett. **74**, 2694 (1995).

[3] J. Kurchan, J. Phys. A: Math. Gen. **31** 3719 (1998).

[4] J. L. Lebowitz and H. Spohn, J. Stat. Phys. **95** 333 (1999).

[5] C. Jarzynski, Phys. Rev. Lett. **78**, 2690 (1997); C. Jarzynski, Phys. Rev. E **56**, 5018 (1997).

[6] G. E. Crooks, J. Stat. Phys. **90**, 1481 (1998); G. E. Crooks, Phys. Rev. E **61**, 2361 (2000).

[7] V. Y Chernyak, M. Chertkov and C. Jarzynski, J. Stat. Mech. P08001 (2006)

[8] R. García-García, D. Domínguez, V. Lecomte, and A. B. Kolton, Phys. Rev. E **82**, 030104(R) (2010)

[9] R. García-García, V. Lecomte, A. B. Kolton, and D. Domínguez, J. Stat. Mech. P02009 (2012)

[10] T. Speck and U. Seifert, J. Stat. Mech. L09002 (2007)

[11] T. Ohkuma and Takao Ohta, J. Stat. Mech. P10010 (2007)

[12] T. Mai and A. Dhar, Phys. Rev. E **75**, 061101 (2007)

[13] C. Aron, G. Biroli, and L. F. Cugliandolo, J. Stat. Mech. P11018 (2010)

[14] F. Zamponi, F. Bonetto, L. F. Cugliandolo, and J. Kurchan, J. Stat. Mech. P09013 (2005)

[15] A. V. Chechkin and R. Klages, J. Stat. Mech. L03002 (2009)

[16] R. García-García, Phys. Rev. E **86**, 031117 (2012)

[17] G. M. Wang, E. M. Sevick, E. Mittag, D. J. Searles, and D. J. Evans, Phys. Rev. Lett. **89**, 050601 (2002)

[18] E. H. Trepagnier, C. Jarzynski, F. Ritort, G. E. Crooks, C. J. Bustamante and J. Liphardt, PNAS **101** 15038 (2004)

[19] C. Bustamante, J. Liphardt, and F. Ritort, Physics Today, **58** 43 (2005).

[20] D. M. Carberry, M. A. B. Baker, G. M. Wang, E. M. Sevick, and D. J. Evans, J. Opt. A.: Pure Appl. Opt. **9**, S204 (2007)

[21] F. Ritort, Advances in Chemical Physics **137**, 31 (2008) Ed. Wiley & Sons. arXiv:0705.0455v1

[22] A. N. Gupta, A. Vincent, K. Neupane, H. Yu, F. Wang, and M. T. Woodside, Nature Physics **7**, 631 (2011)

[23] I. Dornic, and C. Godrèche, J. Phys. A: Math. Gen. **31**, 5413 (1998)

[24] T.J. Newman, and Z. Toroczkai, Phys. Rev. E **58**, R2685 (1998)

[25] Takahiro Hatano and Shin-ichi Sasa, Phys. Rev. Lett. **86**, 3463 (2001)

[26] G. E. Crooks and C. Jarzynski, Phys. Rev. E **75**, 021116 (2007)

[27] J. Prost, J.-F. Joanny and J. M. R. Parrondo, Phys. Rev. Lett. **103** 090601 (2009)

[28] N. Goldenfeld, *Lectures on phase transitions and the renormalization group* (Perseus Books Publishing, L. L. C., 1992)